




Laura Bernard and
Alexandre Le Tiec
Paris Observatory



Gravitational waves

The Universe in my pocket

Answer on overleaf



Which of these
objects do not
emit gravitational
waves?




Quiz

Current detectors

Existing gravitational-wave detectors are based on the principle of **optical interferometry**: they measure tiny variations in length by superimposing laser light on itself.

- They consist of two perpendicular arms, each about a kilometre long, through which light travels. When a gravitational wave passes through them, the difference in length between the two arms varies slightly. This variation is of the order of one thousandth of the size of an atomic nucleus, or 0.0000000000000001 metres.
- Four detectors are in operation: the two **LIGO** instruments in the United States (at Livingston and Hanford);
- the Franco-Italian **Virgo** observatory, near Pisa in Italy;
- the **KAGRA** detector in Japan.




Operating diagram of an interferometric gravitational-wave detector.


Multi-messenger astronomy

13

In August 2017, the coalescence of two neutron stars was observed for the first time. Almost simultaneously, **LIGO** and **Virgo** measured the gravitational wave signal emitted as the two compact bodies spiralled, and the **FERMI** satellite detected the gamma-ray burst (see TUIMP No. 9) resulting from their merger. In the days that followed, numerous telescopes observed the various electromagnetic counterparts (visible, infrared, radio, etc.) of this event. This historic observation inaugurated what is known as **multi-messenger** astronomy, with not only electromagnetic waves being detected but also gravitational waves and high-energy particles. It led to a number of breakthroughs, confirming the propagation of gravitational waves at the speed of light, the suspected link between short gamma-ray bursts and the coalescence of neutron stars, and the mechanism of gold synthesis.



The gamma-ray signal observed by FERMI and the position of the source predicted by LIGO-Virgo (in green) [Credits: LIGO-Virgo, FERMI].




Signals from the merger of two neutron stars: top, gamma ray; bottom, increase in the frequency of the gravitational wave. [Credits: LIGO-Virgo, FERMI]

4




Artist's view of a Type Ia supernova [Credit: ESO].



Artist's view of an isolated neutron star [Credit: NASA].



Artist's depiction of a pair of white dwarf stars [Credit: ESO].



Representation of a pair of neutron stars [Credit: R. Hurt/Calttech-JPL]

Local sources


The main sources of gravitational waves are massive, compact stars (see TUIMP No. 9), such as black holes, neutron stars and white dwarfs, either singly or in pairs. The following localised sources can be distinguished:

- **Binary systems of compact objects** such as spiralling and merging black holes or neutron stars, either galactic or extra-galactic;
- **Binary white dwarf stars** in the Milky Way;
- **Isolated**, slightly asymmetric, rotating **neutron stars** in the galactic neighbourhood;
- **Explosion of massive stars** (supernovae) in our galaxy, leading to the formation of neutron stars or black holes.

Gravitational waves are **small vibrations** in the **structure of space-time** that propagate at the speed of light. They are transverse waves which means that the displacement in space-time is perpendicular to the direction of propagation. They were predicted by **general relativity**, the theory of gravitation formulated by Albert Einstein in 1915.


The first indirect evidence of their existence was the observation by Hulse and Taylor in 1974 of their effect on the orbital period of a pair of neutron stars. The first **instrumental detection** of a gravitational wave took place in 2015 using the **LIGO** detectors. This gravitational wave came from the coalescence of two black holes of about thirty solar masses each.

What are they?



[Credit: D. J. Champion]

Depiction of an array of pulsars. Each line of sight to a pulsar acts as an arm of the interferometer on which the passage of a gravitational wave is measured.



[Credit: NASA]

Representation of the expansion of the Universe from the period of inflation to the present day.

Translation: Stean Kurtz
TUIMP Creative Commons



To find out more about this collection and the themes visit <http://www.tuimp.org>.




Cover image: Numerical simulation of a pair of black holes and visualisation of the gravitational waves generated when they merge [credit: Michael Kopitz/Albert Einstein Institute].

The Universe in my pocket N° 18
This mini-book was written by Laura Bernard and Alexandre Le Tiec from Paris Observatory (France).

Representation of a binary system of black holes
MPI for Gravitational Physics / Institute for Theoretical Physics, Frankfurt / Zuse Institute Berlin



Artist's view of a neutron star




The Crab Nebula, a supernova remnant.
Credit: HST

All these objects are (or have been) sources of gravitational waves, with the exception of the planetary nebula.

Response



Artist's impression of a supernova




The planetary nebula IC 418
Credit: HST


Future gravity wave detectors

Following on from existing detectors, the European Union plans to build the **Einstein Telescope**, a new gravitational wave observatory. It will have a triangular configuration, greater vibration isolation because it will be placed underground, and cryogenic cooling technology for the mirrors.

The European Space Agency is developing **LISA**, a space-based gravitational-wave detector, to eliminate terrestrial disturbances such as seismic noise. It will consist of three satellites several million kilometres apart, enabling it to observe particularly massive sources, complementary to those seen from Earth.



[Credit: Marco Kraan / Nikhef]



The future LISA space detector. The three satellites in triangular configuration follow the Earth's orbit.


Diffuse sources

When gravitational waves generated by a very large number of localised sources are superimposed, they can no longer be distinguished from one another. The result is a **stochastic astrophysical background**.

In addition, various more speculative physical phenomena produced shortly after the Big Bang (see TUIMP n°12) might generate a **cosmological stochastic background**. These include :

- **cosmic strings**, resulting from a sudden change of state in the energy and material content of the primordial Universe;
- **primordial black holes**, which are thought to have formed as a result of large fluctuations in the energy density of the early Universe;
- **inflation**, a period of rapid cosmic expansion that took place a fraction of a second after the Big Bang.

[Credit: NASA]



An artistic depiction of two black holes orbiting each other under the effect of their mutual gravitational attraction. Their orbital motion generates gravitational waves.