


The planetary nebula Hb 12, ionised by a 4.5 000 K star. The strongest lines in its spectrum are the hydrogen recombination lines H $\alpha$  and H $\beta$  and the forbidden lines of the O $^{++}$  ion (oxygen atom that has lost two electrons), noted as [O III].

## The Universe in my pocket



Grażyna Stasińska  
Paris Observatory

**Spectra of ionised nebulae**

Nebulae are clouds of diffuse gas. They can be **ionised** by young massive stars with temperatures around 40000K (these are the H II regions) or by less massive evolved stars that can exceed 100000K (these are the planetary nebulae).

The spectra of **ionised** nebulae are very different from stellar spectra. While the latter show mostly **absorption** lines, the bulk of the light in nebulae is **emitted** in only a few lines, which originate either from **recombinations** of hydrogen and helium or from **collisions** with free electrons in the gas.


These **collisional** lines are not observed in stars and were first attributed to an unknown element, called 'nebulium'. It was not until 1928 that Ira Bowen showed that these lines originate from known elements but only occur at very low densities. They are called '**forbidden lines**'.





Which of these objects has emission lines in its spectrum?



Answer on the back



## Quiz




**Quasar spectra**

Quasars are objects located at very large distances and contain a supermassive black hole at their centre, which attracts the surrounding matter (see TUIMP 6). Before falling into the black hole, the matter is coiled into an 'accretion disc' and heated to hundreds of thousands of degrees. This results in a very blue spectrum.

The **emission** lines are broadened and redshifted (this shift is called 'redshift').

The broadening and redshift are due to the Doppler effect (see TUIMP 15), which changes the frequency of the light wave as a function of the speed of the source relative to the observer. The lines are redshifted due to the recession of quasars due to the expansion of the Universe and broadened due to the rotation of matter around the black hole.


A quasar located at a redshift  $z$  of 0.548, which corresponds to a distance of about 10 billion light years.




**The temperatures of the stars**

Not all stars have the same colour. The cooler ones are red. The hotter ones are blue. The Sun, with a surface temperature of 5500°C (5800 K), is yellow. These colour differences arise because of the way the shape of a star's radiation spectrum varies with temperature, as shown in the figure on the opposite page.

Below are visible spectra of real stars of different types (O, B, A, F, G, K, M). Each type has its own temperature. In addition to the overall distribution of radiation intensity, there are also **absorption** lines of varying depth due to elements present in the atmosphere of the stars in the form of atoms or ions.



Radiation intensity of a hot body as a function of wavelength for various values of the temperature.



Spectra of stars of different types in the visible light range of the electromagnetic spectrum.


In 1835 the French philosopher Auguste Comte said that we would never know what the stars are made of.

However, Isaac Newton had already shown that if a beam of light from the Sun is spread apart by a prism, a spot with the colours of the rainbows is obtained: a **'spectrum'** (see TUIIMP 2).


In 1814, Joseph von Fraunhofer built a spectrograph that discovered more than 500 dark **lines** in the spectrum of the Sun. But it was not until 1860 that Gustav Kirchhoff showed that these lines originate from chemical elements in the upper layers of the Sun. The identification of these lines began soon afterwards, contradicting Comte's pessimistic prediction.

**This was the beginning of astrophysics**, the branch of astronomy that studies the nature of stars by analysing the radiation they emit.

Here are the spectra of two stars that differ only in their chemical composition.



Above: a G-type star of normal composition (similar to the Sun).



Above: a G-type star with a surface very rich in carbon (produced in the star's interior and brought to the surface by convection). The deep, broad lines and the depression at shorter wavelengths are due to carbonaceous molecules.


## The spectra of galaxies

A galaxy contains millions or even billions of stars and sometimes gas. The spectrum of a galaxy is therefore expected to look like a combination of stellar spectra and possibly nebular spectra.


Elliptical galaxies (see TUIIMP 3 and 23) contain no gas and no new stars have formed in them for a long time. They are populated only by old, red stars. All the massive stars they once contained have exploded as supernovas. Their spectra show only **absorption lines**.

Spiral galaxies contain gas and massive stars (**O-** and **B-type**) capable of **ionising** the gas. Thus their spectra show intense **emission** lines, superimposed on a spectrum dominated by hot stars.


An elliptical galaxy. Its spectrum, obtained as part of the Sloan Digital Survey (SDSS), shows decreasing intensity towards shorter wavelengths because most of the stars of the galaxy are red. The characteristic **absorption** lines of these stars can be seen.







A spiral galaxy. Its spectrum, showing **emission** lines, is similar to that of an HII region.



All other objects have **stellar-like** spectra.



Only the planetary nebula has **emission lines** in its spectrum.



## The composition of the stars

Today, thanks to the **absorption** lines observed in the spectra of stars, astronomers know which elements are present in their atmospheres and can measure their abundance.

The chemical composition of the atmosphere of a star is, in general, identical to that of the molecular cloud in which it formed. The interior of the star has a different chemical composition from the atmosphere because of the nucleosynthetic reactions that occur there (see TUIIMP 14) but this is not measured directly.

It can be seen that, broadly speaking, stars have a chemical composition similar to that of the Sun. However, stars in the outermost parts of the Galaxy tend to be less rich in elements heavier than helium, because they are less enriched by nucleosynthesis products.